

AD-A254 370




AD

TECHNICAL REPORT ARCCB-TR-92037

## MULTIFRACTAL ANALYSIS AT NEGATIVE $q$

MARK JOHNSON  
L. V. MEISEL



AUGUST 1992



US ARMY ARMAMENT RESEARCH,  
DEVELOPMENT AND ENGINEERING CENTER  
CLOSE COMBAT ARMAMENTS CENTER  
BENÉT LABORATORIES  
WATERVLIET, N.Y. 12189-4050



APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED

92 8 19 55

92-23120

#### DISCLAIMER

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

The use of trade name(s) and/or manufacturer(s) does not constitute an official indorsement or approval.

#### DESTRUCTION NOTICE

For classified documents, follow the procedures in DoD 5200.22-M, Industrial Security Manual, Section II-19 or DoD 5200.1-R, Information Security Program Regulation, Chapter IX.

For unclassified, limited documents, destroy by any method that will prevent disclosure of contents or reconstruction of the document.

For unclassified, unlimited documents, destroy when the report is no longer needed. Do not return it to the originator.

# REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE August 1992		3. REPORT TYPE AND DATES COVERED Final	
4. TITLE AND SUBTITLE MULTIFRACTAL ANALYSIS AT NEGATIVE $q$				5. FUNDING NUMBERS AMCMS No. 6111.01.91A1.1	
6. AUTHOR(S) Mark Johnson and L. V. Meisel					
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Army ARDEC Benet Laboratories, SMCAR-CCB-TL Watervliet, NY 12189-4050				8. PERFORMING ORGANIZATION REPORT NUMBER ARCCB-TR-92037	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army ARDEC Close Combat Armaments Center Picatinny Arsenal, NJ 07806-5000				10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES Submitted to <i>Physical Review Letters</i> .					
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited.				12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words)  A box-based correlation integral algorithm for the determination of generalized fractal dimensions is described. The algorithm has been tested for Euclidean curves, Koch symmetric and asymmetric triadic snowflakes, split snowflake halls, and the multifractal construction based on Mandelbrot's 13-element generator. The technique is well-suited to the analysis of experimental data with inherent uncertainty, it is efficient, and it converges to analytic results for the model point sets at all $q$ values.					
14. SUBJECT TERMS Fractal, Multifractal, Box-Counting, Algorithm				15. NUMBER OF PAGES 7	
				16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT UL		

## TABLE OF CONTENTS

	<u>Page</u>
INTRODUCTION .....	1
THEORY .....	1
RESULTS AND CONCLUSIONS .....	2
REFERENCES .....	4

## LIST OF ILLUSTRATIONS

1. Analytic and BBCI measurements of the fractal dimension  $D(q)$  versus level for randomly-oriented split snowflake halls ..... 5
2. Analytic and BBCI values of the fractal dimension  $D(q)$  versus  $q$  for level 8 split snowflake halls ..... 6

Accession For	
NTIS CRA&I	✓
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

DTIC QUALITY INSPECTED 5

## INTRODUCTION

Block, von Bloh, and Schellnhuber (BBS) (ref 1) presented a box-counting algorithm for the measurement of generalized (fractal) dimensions, which they refer to as "Efficient Box-Counting" (EBC). Meisel, Johnson, and Cote (ref 2) described a box-counting algorithm, which is particularly suited to the analysis of experimental data with inherent uncertainty and large data sets, which they referred to as "Agglomeration Box-Counting" (ABC). Atmanspacher, Scheingraber, and Wiedenmann (ref 3) applied techniques similar in spirit to EBC and ABC, which they referred to as "histogram techniques," and a technique based on the generalized correlation integral (ref 4), to the determination of the multifractal properties of the distribution of galaxies on the celestial sphere (ref 3).

It was shown in Reference 2 that although box-counting algorithms converge to analytic results for model fractal point sets at  $q \geq 0$ , they fail to converge to analytic results at  $q < 0$ . Similarly, Atmanspacher et al. (ref 3) reported consistent results for the left branch of the  $f(\alpha)$  spectrum ( $q \geq 0$ ), but noted discrepancies for the right-hand branch ( $q < 0$ ). It is evident that a reliable algorithm for the determination of  $D(q)$  at  $q < 0$  or equivalently for the determination of the right-hand branch of the  $f(\alpha)$  spectrum is urgently required.

The present work describes a Box-Based Correlation Integral (BBCI) method for multifractal analysis. The BBCI algorithm has been tested for a selection of model multifractal point sets, which are subject to simple analysis along the lines in Halsey et al. (ref 5). BBCI values of  $D(q)$  converge close to analytic values at all  $q \in [-25, 25]$  for the model multifractal point sets. The algorithm is well-suited to experimental data application, and it converges at least as rapidly as ABC at  $q > 1$ .

## THEORY

The Box-Based Correlation Integral (BBCI) algorithm is implemented as follows:

1. Define a set of nested hypercubes ("boxes") appropriate for the given (or anticipated) point set  $S$ . We refer to these boxes as elementary hypercubes or elementary boxes. For experimental data, the appropriate choice of elementary box edge  $E_0$  should reflect the inherent uncertainty of the coordinates of the point set within limits set by storage requirements.

Storage requirements are effectively determined by the choice of elementary hypercubes rather than the size of the point set. Thus, large point sets can be accommodated with relatively modest resources.

2. Compute or measure the occupation numbers  $n_i$  for each of the elementary hypercubes. Define  $N$  as

$$N = \sum_{i=1}^{\text{elementary boxes}} n_i.$$

In our tests  $N$  ranged between about  $10^3$  and  $10^9$ .

3. Define a reference set that comprises a subset of the occupied elementary boxes. Let  $N_{\text{ref}}$  be the number of elementary boxes in the reference set. In our tests with  $N = 768^3$  elementary boxes, we took  $n_{\text{ref}}$  as the smaller of the total number of occupied boxes or 15,000.

4. For each member of the reference set, define a set of hypercube edge lengths,  $E = (2n+1) E_0$ , where  $n=0,1,2,\dots,n_{\text{MAX}}$ . In our tests with  $N = 768^3$  elementary boxes, we took  $n_{\text{MAX}} = 24$ .

5. For each  $q$  of interest:

a. Compute the box-based generalized correlation integrals  $C(q, E, E_0)$ , which are defined as

$$C(q, E, E_0) = \left\{ \frac{1}{N_2} \sum_r^{N_{ref}} \left[ \frac{1}{N} \left( \sum_j^{\text{elementary boxes}} n_r n_j G(E, E_0, \mathbf{x}_r - \mathbf{x}_j) - 1 \right) \right]^{q-1} \right\}^{\frac{1}{q-1}}$$

for the  $E$  values defined in step 4, where  $r$  runs over the reference set,  $j$  runs over all elementary boxes,

$$N_2 = \sum_r^{N_{ref}} n_r$$

and the function

$$G(E, E_0, \mathbf{x}) = 1, \text{ if } |\mathbf{x}_i| \leq (E - E_0)/2, \text{ for all components of } \mathbf{x} \\ = 0, \text{ otherwise,}$$

selects the elementary boxes contained in larger hypercubes of edge  $E$ . The vectors  $\mathbf{x}_r$  and  $\mathbf{x}_j$  point to elementary hypercubes rather than members of the point set. Reducing the sum over elementary boxes in the side  $E$  hypercube by unity eliminates self-correlation contributions.

The box-based generalized correlation integral reduces to the standard generalized correlation integral (ref 4) in the case when the coordinates of the members of the multifractal subset in question are precisely known, the elementary hypercubes are small enough to contain at most one element of the multifractal subset, and the covering set is comprised of hypercubes. That is,

$$C(q, E) = \lim_{E_0 \rightarrow 0^+} C(q, E, E_0).$$

Computation was slowed down and convergence was not improved by employing (approximate) hyperspheres rather than hypercubes. Also note that special techniques are required near  $q = 1$ ; here, we restricted  $q$  such that  $|q - 1| > 1/2$ .

b. Obtain  $D(q)$  and the corresponding rms error in  $D(q)$  by linear regression on  $\ln[C(q, E, E_0)] = \text{const} + D(q) \ln[E]$  for  $E \in \{(2n+1) E_0 | n=1, 2, \dots, n_{MAX}\}$ .

## RESULTS AND CONCLUSIONS

The BBCI algorithm has been tested for Euclidean curves, Koch symmetric and asymmetric triadic snowflakes, split snowflake halls, and the multifractal construction based on Mandelbrot's 13-element generator. The  $D(q)$  determined by BBCI converged to values near the analytic  $D(q)$  values for  $q$  ranging between -25 and +25 in all cases as  $N$  was increased.

Figure 1 shows results of application of the BBCI algorithm to split snowflake halls at five values of  $q$ . The algorithm was applied to a randomly-oriented construction centered on 768 by 768 square boxes;  $N_{\text{tot}}$  was the lesser of 15,000 or the total number of occupied boxes. The horizontal lines are the analytic values. Convergence was within about 2 percent for  $q < 1$  and within 1 percent for  $q > 1$  in this test. Note that convergence is about an order of magnitude faster for  $q > 1$  than it is for  $q < 1$ . Except that convergence is faster for constructions with lower  $D(0)$ , Figure 1 is typical of convergence obtained by application of the BBCI algorithm to the constructions tested.

Figure 2 shows the results of application of the BBCI algorithm as in Figure 1 for  $N = 2.14 \cdot 10^4$  and analytic values of  $D(q)$  versus  $q$  for split snowflake halls. The increase in the error in  $D(q)$  with  $|q|$  and the order of the discrepancies are apparent.

The rms errors found in the linear regressions were of the order of the differences between analytic and BBCI results for  $D(q)$ . An expanded discussion of the results of application of BBCI to model fractals and of the effects of variations of  $N_{\text{tot}}$ , number of elementary boxes, and  $n_{\text{MAX}}$  will be presented in a future publication.

## REFERENCES

1. A. Block, W. von Bloh, and H.J. Schellnhuber, *Phys. Rev. A*, Vol. 42, 1990, p. 1869. Other recent references to sorting algorithms for the determination of  $D(0)$  include: L.S. Liebovitch and T. Toth, *Phys. Lett. A*, Vol. 141, 1989, p. 386; X.J. Hou, R.G. Gilmore, G.B. Mindlin, and H.G. Solari, *Phys. Lett. A*, Vol. 151, 1990, p. 43.
2. L.V. Meisel, M. Johnson, and P.J. Cote, *Phys. Rev. A*, Vol. 45, 1992, p. 6989.
3. H. Atmanspacher, H. Scheingraber, and G. Wiedenmann, *Phys. Rev. A*, Vol. 40, 1989, p. 3954.
4. For the standard generalized correlation integral see, G. Paladin and A. Vulpiano, *Lett. Nuovo Cimento*, Vol. 41, 1984, p. 82 and K. Pawelzik and H.G. Schuster, *Phys. Rev. A*, Vol.35, 1987, p. 481.
5. Thomas C. Halsey, Mogens H. Jensen, Leo P. Kadanoff, Itamar Procaccia, and Boris I. Shraiman, *Phys. Rev. A*, Vol. 33, 1986, p. 1141.
6. B. B. Mandelbrot, *Fractal Geometry of Nature*, Freeman, New York, 1983.






Figure 1. Analytic and BBCI measurements of the fractal dimension  $D(q)$  versus level for randomly-oriented split snowflake halls. (The number of points in the construction  $N = 11^{\text{level}}$ .)




Figure 2. Analytic and BBCI values of the fractal dimension  $D(q)$  versus  $q$  for level 8 split snowflake halls. The line is the analytic curve.

# TECHNICAL REPORT INTERNAL DISTRIBUTION LIST

	NO. OF COPIES
CHIEF, DEVELOPMENT ENGINEERING DIVISION	
ATTN: SMCAR-CCB-DA	1
-DC	1
-DI	1
-DR	1
-DS (SYSTEMS)	1
CHIEF, ENGINEERING SUPPORT DIVISION	
ATTN: SMCAR-CCB-S	1
-SD	1
-SE	1
CHIEF, RESEARCH DIVISION	
ATTN: SMCAR-CCB-R	2
-RA	1
-RE	1
-RM	1
-RP	1
-RT	1
TECHNICAL LIBRARY	5
ATTN: SMCAR-CCB-TL	
TECHNICAL PUBLICATIONS & EDITING SECTION	3
ATTN: SMCAR-CCB-TL	
OPERATIONS DIRECTORATE	1
ATTN: SMCWV-ODP-P	
DIRECTOR, PROCUREMENT DIRECTORATE	1
ATTN: SMCWV-PP	
DIRECTOR, PRODUCT ASSURANCE DIRECTORATE	1
ATTN: SMCWV-QA	

NOTE: PLEASE NOTIFY DIRECTOR, BENET LABORATORIES, ATTN: SMCAR-CCB-TL, OF ANY ADDRESS CHANGES.

# TECHNICAL REPORT EXTERNAL DISTRIBUTION LIST

	<u>NO. OF COPIES</u>		<u>NO. OF COPIES</u>
ASST SEC OF THE ARMY RESEARCH AND DEVELOPMENT ATTN: DEPT FOR SCI AND TECH THE PENTAGON WASHINGTON, D.C. 20310-0103	1	COMMANDER ROCK ISLAND ARSENAL ATTN: SMCRI-ENM ROCK ISLAND, IL 61299-5000	1
ADMINISTRATOR DEFENSE TECHNICAL INFO CENTER ATTN: DTIC-FDAC CAMERON STATION ALEXANDRIA, VA 22304-6145	12	DIRECTOR US ARMY INDUSTRIAL BASE ENGR ACTV ATTN: AMXIB-P ROCK ISLAND, IL 61299-7260	1
COMMANDER US ARMY ARDEC ATTN: SMCAR-AEE	1	COMMANDER US ARMY TANK-AUTMV R&D COMMAND ATTN: AMSTA-DDL (TECH LIB) WARREN, MI 48397-5000	1
SMCAR-AES, BLDG. 321	1	COMMANDER US MILITARY ACADEMY	1
SMCAR-AET-O, BLDG. 351N	1	ATTN: DEPARTMENT OF MECHANICS WEST POINT, NY 10996-1792	
SMCAR-CC	1		
SMCAR-CCP-A	1		
SMCAR-FSA	1	US ARMY MISSILE COMMAND	
SMCAR-FSM-E	1	REDSTONE SCIENTIFIC INFO CTR	2
SMCAR-FSS-D, BLDG. 94	1	ATTN: DOCUMENTS SECT, BLDG. 4484	
SMCAR-IMI-I (STINFO) BLDG. 59	2	REDSTONE ARSENAL, AL 35898-5241	
PICATINNY ARSENAL, NJ 07806-5000			
DIRECTOR US ARMY BALLISTIC RESEARCH LABORATORY ATTN: SLCBR-DD-T, BLDG. 305 ABERDEEN PROVING GROUND, MD 21005-5066	1	COMMANDER US ARMY FGN SCIENCE AND TECH CTR ATTN: DRXST-SD 220 7TH STREET, N.E. CHARLOTTESVILLE, VA 22901	1
DIRECTOR US ARMY MATERIEL SYSTEMS ANALYSIS ACTV ATTN: AMXSY-MP ABERDEEN PROVING GROUND, MD 21005-5071	1	COMMANDER US ARMY LABCOM MATERIALS TECHNOLOGY LAB ATTN: SLCMT-IML (TECH LIB) WATERTOWN, MA 02172-0001	2
COMMANDER HQ, AMCCOM ATTN: AMSMC-IMP-L ROCK ISLAND, IL 61299-6000	1		

NOTE: PLEASE NOTIFY COMMANDER, ARMAMENT RESEARCH, DEVELOPMENT, AND ENGINEERING CENTER, US ARMY AMCCOM, ATTN: BENET LABORATORIES, SMCAR-CCB-TL, WATERVLIET, NY 12189-4050, OF ANY ADDRESS CHANGES.

# TECHNICAL REPORT EXTERNAL DISTRIBUTION LIST (CONT'D)

	<u>NO. OF COPIES</u>		<u>NO. OF COPIES</u>
COMMANDER US ARMY LABCOM, ISA ATTN: SLCIS-IM-TL 2800 POWDER MILL ROAD ADELPHI, MD 20783-1145	1	COMMANDER AIR FORCE ARMAMENT LABORATORY ATTN: AFATL/MN EGLIN AFB, FL 32542-5434	1
COMMANDER US ARMY RESEARCH OFFICE ATTN: CHIEF, IPO P.O. BOX 12211 RESEARCH TRIANGLE PARK, NC 27709-2211	1	COMMANDER AIR FORCE ARMAMENT LABORATORY ATTN: AFATL/MNF EGLIN AFB, FL 32542-5434	1
DIRECTOR US NAVAL RESEARCH LAB ATTN: MATERIALS SCI & TECH DIVISION CODE 26-27 (DOC LIB) WASHINGTON, D.C. 20375	1 1	MIAC/CINDAS PURDUE UNIVERSITY 2595 YEAGER ROAD WEST LAFAYETTE, IN 47905	1
DIRECTOR US ARMY BALLISTIC RESEARCH LABORATORY ATTN: SLCBR-IB-M (DR. BRUCE BURNS) ABERDEEN PROVING GROUND, MD 21005-5066	1		

NOTE: PLEASE NOTIFY COMMANDER, ARMAMENT RESEARCH, DEVELOPMENT, AND ENGINEERING CENTER, US ARMY AMCCOM, ATTN: BENET LABORATORIES, SMCAR-CCB-TL, WATERVLIET, NY 12189-4050, OF ANY ADDRESS CHANGES.